arXiv:cond-mat/9710275v1 24 Oct 1997

The Hydrodynamic Equations of Superfluid Mixtures in Magnetic Traps

Tin-Lun Ho and V.B. Shenoy
Physics Department, The Ohio State University, Columbus, Ohio 43210

(February 1, 2008)

Abstract

The hydrodynamics equations of binary mixtures of Bose gases, Fermi gases, and mixtures of Bose and Fermi gases in the presence of external potentials are derived. These equations can be applied to current experiments on mixtures of atomic gases confined in magnetic traps.

The recent discoveries of Bose-Einstein condensation of atomic gases [1–3] have generated much interest in studying dilute Bose gases trapped in external potentials. While most of the experiments are performed on trapped atoms of the same kind, the current traps can also hold different kinds of atoms at the same time, providing opportunities to study mixtures of Bose gases, mixtures of Fermi gases and mixtures of Bose and Fermi gases. In a recent paper, we have shown [4] that there is a great variety of ground state structures in the binary mixtures of Bose gases. These structures can be experimentally accessed by varying the number of trapped Bosons of each kind. The two Bosons can be different alkalis, different isotopes of the same alkali, or different hyperfine spin (F) states of the same alkali isotope. Recently, a mixture of different hyperfine states of ^{87}Rb ($F = 1$ and $F = 2$) has been produced by the JILA group [5], creating a system of interpenetrating Bose superfluids for the first time.

In the case of superfluid ^{4}He, its superfluid properties are contained in the well known two-fluid hydrodynamic equations of Landau [6] that incorporate the effect of broken gauge symmetry into hydrodynamics. In order to understand the properties of mixtures, it is important to derive the corresponding hydrodynamic equations, which is the goal of this paper. Our derivation is a standard one. It follows from general principles such as conservation laws, broken symmetries, thermodynamic relations and the second law of thermodynamics [6,7]. It will be clear from the derivation that the form of the hydrodynamic equations is independent of the density of the gases, as long as the symmetry of the system remains unchanged. What distinguishes a dilute and a dense system is the explicit form of the coefficients entering these equations, which are related to the thermodynamic potential of the system. In this paper, we will focus only on the derivation of the hydrodynamic equations of mixtures. The solutions of these equations require the evaluation of the thermodynamic potentials which will be discussed elsewhere. (We have recently solved the hydrodynamic equations of single component dilute Bose gases in spherical traps and have obtained all the first and second sound modes (classified by radial and angular quantum numbers) as a function of temperature [8]).

Before proceeding, it is necessary to comment on the broken symmetry of the alkali gases in magnetic traps. Since the alkali atoms carry spin, the order parameter in the Bose condensed phase should be a spinor field instead of a scalar field. However, in a magnetic trap, only the spin states locally aligned with the magnetic field $\mathbf{B}(\mathbf{r})$ are trapped [9]. The resulting quantum gas therefore has the spin field "frozen" along the local field, leaving particle motion as the only degree of freedom. As a result, the spin field reduces to a scalar. On the other hand, even though the spin field is frozen, the spinor nature is not completely erased. In a recent paper [9], we have shown that the spatial variations of the spin field will give rise to a Berry phase which can lead to a number of dramatic effects. While the Berry phase effects in current traps are small, they can be magnified by changing the geometry of the trap. For simplicity, we shall only consider those traps where the Berry phase effects are negligible. One should, however, keep in mind that the hydrodynamics equations derived here are limited to traps of this kind.

Since both scalar Bose superfluids and s-wave Fermion superfluids have identical broken $U(1)$ gauge symmetry, and since the hydrodynamics of a system depends entirely on the nature of its broken symmetry, the hydrodynamic equations of these two systems are identical. The hydrodynamics of binary mixtures of quantum gases (which can be either Fermions or Bosons) therefore fall in three categories, namely, mixtures of superfluids, mixtures consisting of a normal fluid and a superfluid, and mixtures of normal fluids. The superfluids can be the scalar condensates of Bosons or the s-wave BCS condensates of Fermions. The normal fluid can be the either Bosons or Fermions in their normal state.

Among all the cases considered, the hydrodynamics of binary superfluid mixtures in a trap is the most complex. We shall focus on this case. As we shall see, once we derive the hydrodynamic equations for this system, extensions to other mixtures is easy. For a binary mixture of Bosons, in general, there will be regions in the trap where the two species coexist and regions where there is no coexistence. Furthermore, in the region of coexistence, each Boson species may be in their normal or superfluid states. Since the hydrodynamic modes in each of different regions are connected to one another, not only does one need to know the hydrodynamic equations of interpenetrating superfluids, but also those of the surrounding mixtures of normal and superfluids.

The organization of this paper is as follows. In Section **I**, we will discuss the hydrodynamic variables of binary mixtures of scalar superfluids and their Galilean transformation properties. The hydrodynamic equations for this system of interpenetrating scalar superfluids will be derived in Section **II**. The explicit form of the hydrodynamic equations (with and without dissipation) will be presented. Section **III** is a summary of the hydrodynamic equations for other mixture systems.

I : The hydrodynamic variables of a binary superfluid mixtures, conservation laws, and their Galilean transformation properties :

Before studying hydrodynamics, it is necessary to study the equilibrium thermodynamics of homogeneous systems, which is described by the entropy function S. S is a function of a set of thermodynamic variables $\{X_i\}$, which are quantities specifying the physical state of the system, $S = S(\{X_i\})$. To study binary mixtures, it is useful to consider the entropy functions of the following sequence of systems with increasing complexity :
(i) Single component normal fluids : In this case, $\{X_i\} = (E, \mathbf{P}, N, V)$, $S = S(E, \mathbf{P}, N, V)$, where $E, \mathbf{P}, N$ and V are the total energy, momentum, particle number and volume of the system respectively. The meaning of $E, \mathbf{P}$ and V will remain unchanged in the following

examples (ii) to (v).

(ii) Binary mixtures of normal fluids : Each component in the mixture can be either a Boson or a Fermion. In this case $S = S(E, \mathbf{P}, N_1, N_2, V)$, where N_1 and N_2 are the particle numbers of the two normal fluids labelled as **1** and **2** respectively. The reason that S is a function of N_1 and N_2 but not a function separate momenta $\mathbf{P}_1$ and $\mathbf{P}_2$ is because while Boson **1** does not turn into Boson **2**, they can exchange momenta through particle collisions. As a result, the equilibrium state of the system is characterised by a single momentum.

(iii) Single component scalar superfluid : This is the well known case of ^{4}He. The system now acquires a broken gauge symmetry $< \hat{\psi}(\mathbf{r}) >= | < \hat{\psi}(\mathbf{r}) > |e^{i\theta(\mathbf{r})}$, where $\hat{\psi}$ is the field operator. In this case, $S = S(E, \mathbf{P}, N, V; \mathbf{v}_s)$, where $\mathbf{v}_s = (\hbar/m)\nabla\theta$ is the superfluid velocity describing the phase winding in the system, and m is the mass of the Boson.

(iv) Binary mixtures consisting of a scalar superfluid and a normal fluid : Denoting the superfluid and the normal fluid components as **1** and **2** respectively, we have for this case $S = S(E, \mathbf{P}, N_1, N_2, V; \mathbf{v}_{s1})$. The reason that S is a function of total momentum $\mathbf{P}$ instead of separate momenta $\mathbf{P}_1$ and $\mathbf{P}_2$ is the same as that for normal fluid mixtures.

(v) Binary mixtures of superfluids : We denote the two superfluids as 1 and 2, and their masses by m_1 and m_2 respectively. From the discussion in cases (i) to (iv) above, we have

$$S = S(E, \mathbf{P}, N_1, N_2, V; \mathbf{v}_{s1}, \mathbf{v}_{s2}), \tag{1}$$

with

$$\mathbf{v}_{s\alpha} \equiv \frac{\hbar}{m_\alpha}\nabla\theta_\alpha, \qquad \alpha = 1, 2. \tag{2}$$

From this point onwards, we shall consider only case (v). The hydrodynamic equations of cases (i) to (iv) can be obtained from the equations derived for case (v) by setting the appropriate superfluid densities (defined below) to zero. (See section **(III)**).

The extensive property of S implies

$$S(E, \mathbf{P}, N_1, N_2, V; \mathbf{v}_{s1}, \mathbf{v}_{s2}) = Vs(\epsilon, \mathbf{g}, \rho_1, \rho_2, \mathbf{v}_{s1}, \mathbf{v}_{s2}), \tag{3}$$

where $s = S/V$, $\epsilon = E/V$, $\mathbf{g} = \mathbf{P}/V$ are the entropy density, the energy density, the momentum density, and $\rho_\alpha = m_\alpha N_\alpha/V$ is the mass density of the species labelled α in the mixture, ($\alpha = 1, 2$). The arguments $\{\epsilon, \mathbf{g}, \rho_\alpha, \mathbf{v}_{s\alpha}\}$ of the entropy density s will be referred to as the "hydrodynamic variables" of the system, $\{x_i\}$. Defining temperature T, chemical potential (per unit mass) μ_α, "normal fluid velocity" $\mathbf{v}_n$, and the vector $\mathbf{b}_\alpha$ as $\partial s/\partial\epsilon \equiv 1/T, \partial s/\partial\rho_\alpha \equiv -\mu_\alpha/T, \partial s/\partial\mathbf{g} \equiv -\mathbf{v}_n/T$, and $\partial s/\partial\mathbf{v}_{s\alpha} \equiv -\mathbf{b}_\alpha/T$ respectively, we can write

$$\mathrm{d}s = \frac{1}{T}\mathrm{d}\epsilon - \frac{\mathbf{v}_n}{T}\cdot\mathrm{d}\mathbf{g} - \sum_{\alpha=1}^{2}\frac{\mu_\alpha}{T}\mathrm{d}\rho_\alpha - \sum_{\alpha=1}^{2}\frac{\mathbf{b}_\alpha}{T}\cdot\mathrm{d}\mathbf{v}_{s\alpha}. \tag{4}$$

Defining the pressure as $P = T(\partial S/\partial V)_{E,\mathbf{P},N}$, and using the extensivity relation given by Eq.(3), we have

$$P = Ts - \epsilon + \mu_1\rho_1 + \mu_2\rho_2 + \mathbf{v}_n\cdot\mathbf{g}, \tag{5}$$

and the Gibbs-Duham relation

$$\mathrm{d}P = s\mathrm{d}T + \sum_{\alpha=1}^{2} \rho_\alpha \mathrm{d}\mu_\alpha + \mathbf{g}\cdot \mathrm{d}\mathbf{v}_n - \sum_\alpha \mathbf{b}_\alpha \cdot \mathrm{d}\mathbf{v}_{s\alpha}, \tag{6}$$

which shows the pressure is a function of the intensive variables $(T, \mu_\alpha, \mathbf{v}_n, \mathbf{v}_{s\alpha})$. In analogy to binary mixtures of classical fluids, it is useful to introduce the concentration variable

$$c = \rho_1/\rho, \quad 1 - c = \rho_2/\rho, \tag{7}$$

where $\rho = \rho_1 + \rho_2$. If we define

$$\mu \equiv \mu_1 c + \mu_2(1-c), \tag{8}$$

then Eq.(4) can be written as

$$T\mathrm{d}s = \mathrm{d}\epsilon - \mathbf{v}_n \cdot \mathrm{d}\mathbf{g} - \mu \mathrm{d}\rho - (\mu_1 - \mu_2)\rho \mathrm{d}c - \sum_\alpha \mathbf{b}_\alpha \cdot \mathrm{d}\mathbf{v}_{s\alpha}. \tag{9}$$

Likewise, Eq.(5) can be written as

$$P = Ts - \epsilon + \mu\rho + \mathbf{v}_n \cdot \mathbf{g}, \tag{10}$$

while the Gibbs-Duham relation can be cast in the form

$$\delta P = s\delta T + \mathbf{g}\cdot \delta\mathbf{v}_n + \rho\delta\mu - (\mu_1 - \mu_2)\rho\delta c - \sum_\alpha \mathbf{b}_\alpha \cdot \delta \mathbf{v}_{s\alpha}. \tag{11}$$

The Galilean transformation properties of the hydrodynamic variables : As we shall see, the derivation of the hydrodynamic equations can be simplified by using the Galilean transformation (GT) properties of the hydrodynamic variables X_i and their derivatives $\partial S/\partial X_i$. A Galilean transformation is a space-time transformation ($\mathbf{r} \to \mathbf{r}' = \mathbf{r} + \mathbf{u}t$, $t \to t' = t$), where $\mathbf{u}$ is the relative velocity of the two inertial frames. Under this transformation, a scalar function F of $\mathbf{r}$ and t will change to F', which is a function of $\mathbf{r}'$ and t'. In particular, we have

$$\rho'_\alpha = \rho_\alpha, \quad \alpha = 1, 2 \tag{12}$$

$$\mathbf{g}' = \mathbf{g} + \rho\mathbf{u}, \quad \text{where} \quad \rho = \rho_1 + \rho_2, \tag{13}$$

$$\epsilon' = \epsilon + \mathbf{g}\cdot\mathbf{u} + \rho u^2/2, \tag{14}$$

and

$$\mathbf{v}'_{s\alpha} = \mathbf{v}_{s\alpha} + \mathbf{u}. \tag{15}$$

While the validity of Eqs.(13) and (14) is easy to see, Eq.(15) follows from the Galilean transformation properties of the field operator of the species α,

$$\hat{\psi}'_\alpha(\mathbf{r}', t') = \hat{\psi}_\alpha(\mathbf{r}, t)\mathrm{exp}\left(\frac{im_\alpha \mathbf{u}\cdot\mathbf{r}}{\hbar} + \frac{im_\alpha u^2 t}{2\hbar}\right). \tag{16}$$

The phase θ_α of $< \hat{\psi}_\alpha(\mathbf{r}, t) >$ transforms as

$$\theta'_\alpha = \theta_\alpha + \frac{m_\alpha \mathbf{u}\cdot\mathbf{r}}{\hbar} + \frac{m_\alpha u^2 t}{2\hbar}, \tag{17}$$

hence Eq.(15), which says that $\mathbf{v}_s$ indeed transforms like a velocity field. Finally, we also note that the momentum density of the species α, written as $\mathbf{g}_\alpha$, transforms as

$$\mathbf{g}_\alpha \to \mathbf{g}'_\alpha = \mathbf{g}_\alpha + \rho_\alpha \mathbf{u}, \qquad \text{and} \qquad \mathbf{g} = \mathbf{g}_1 + \mathbf{g}_2. \tag{18}$$

Since the entropy involves counting of states, it is a Galilean invariant; we then have

$$s'\left((\epsilon + \mathbf{g}\cdot\mathbf{u} + \frac{1}{2}\rho u^2), (\mathbf{g} + \rho\mathbf{u}), \rho_1, \rho_2, (\mathbf{v}_{s1} + \mathbf{u}), (\mathbf{v}_{s2} + \mathbf{u})\right) = s\left(\epsilon, \mathbf{g}, \rho_1, \rho_2, \mathbf{v}_{s1}, \mathbf{v}_{s2}\right). \tag{19}$$

Differentiating both sides of Eq.(19) with respect to ϵ and $\mathbf{v}_{s\alpha}$, we find that both T and $\mathbf{b}_\alpha$ are Galilean invariant,

$$T' = T, \quad \mathbf{b}'_\alpha = \mathbf{b}_\alpha. \tag{20}$$

Differentiating both sides of Eq.(19) with respect to $\mathbf{g}$ and ρ_α, we get

$$\mathbf{v}'_n = \mathbf{v}_n + \mathbf{u}, \quad \text{and} \quad \mu'_\alpha = \mu_\alpha - \mathbf{v}_n\cdot\mathbf{u} - \frac{1}{2}u^2, \qquad \alpha = 1, 2. \tag{21}$$

This shows that $\mathbf{v}_n$ also transforms like a velocity field. Finally, differentiating Eq.(19) with respect to $\mathbf{u}$ at $\mathbf{u} = 0$ gives

$$\mathbf{b}_1 + \mathbf{b}_2 = \mathbf{g} - \rho\mathbf{v}_n \equiv \mathbf{g}^0, \tag{22}$$

where the superscript ^{0} refers to quantities evaluated in the $\mathbf{v}_n = 0$ frame. From Eq.(21), we then have

$$\mu_\alpha = \mu^0_\alpha - \frac{1}{2}\mathbf{v}^2_n, \qquad \alpha = 1, 2, \tag{23}$$

and from Eqs.(13), (14), and (21), it is easy to see that the pressure P is a Galilean invariant and can also be written as

$$P = Ts - \epsilon^0 + \mu^0_1\rho_1 + \mu^0_2\rho_2. \tag{24}$$

Rotational symmetry : Under an infinitesimal rotation $\mathbf{a}$, a vector such as $\mathbf{g}$ changes to $\mathbf{g} + \mathbf{a}\times\mathbf{g}$. Since entropy of the system is invariant under such a rotation, we have

$$s\left(\epsilon, (\mathbf{g} + \mathbf{a}\times\mathbf{g}), \rho_\alpha, (\mathbf{v}_{s1} + \mathbf{a}\times\mathbf{v}_{s1}), (\mathbf{v}_{s2} + \mathbf{a}\times\mathbf{v}_{s2})\right) = s(\epsilon, \mathbf{g}, \rho_\alpha, \mathbf{v}_{s1}, \mathbf{v}_{s2}). \tag{25}$$

Differentiating both sides with respect to $\mathbf{a}$, we have

$$\mathbf{v}_n\times\mathbf{g} + \mathbf{b}_1\times\mathbf{v}_{s1} + \mathbf{b}_2\times\mathbf{v}_{s2} = 0, \tag{26}$$

which implies $\mathbf{b}_1 \times \mathbf{v}^0_{s1} + \mathbf{b}_2 \times \mathbf{v}^0_{s2} = 0$. The general form of $\mathbf{b}_\alpha$ satisfying Eq.(26) can be written as

$$\mathbf{b}_1 = \rho_{s1}\mathbf{v}^0_{s1} + \tilde{\rho}(\mathbf{v}^0_{s1} - \mathbf{v}^0_{s2}), \quad \mathbf{b}_2 = \rho_{s2}\mathbf{v}^0_{s2} + \tilde{\rho}(\mathbf{v}^0_{s2} - \mathbf{v}^0_{s1}), \tag{27}$$

where ρ_{s1}, ρ_{s2}, and $\tilde{\rho}$ are parameters which will be referred to as superfluid densities. With this parameterization, using Eq.(22), we have

$$\mathbf{g}^0 = \rho_{s1}\mathbf{v}^0_{s1} + \rho_{s2}\mathbf{v}^0_{s2} \ , \tag{28}$$

$$\mathbf{g} = \rho_n\mathbf{v}_n + \rho_{s1}\mathbf{v}_{s1} + \rho_{s2}\mathbf{v}_{s2}, \qquad \rho_n \equiv \rho - \rho_{s1} - \rho_{s2}, \tag{29}$$

where ρ_n will be refered to as the normal fluid density. For later discussions, it is convenient to define the "diffusion" current $\mathbf{J}^D$ as

$$\mathbf{g}^0_1 \equiv \mathbf{b}_1 + \mathbf{J}^D, \qquad \mathbf{g}^0_2 \equiv \mathbf{b}_2 - \mathbf{J}^D. \tag{30}$$

Using the above definition of $\mathbf{J}^D$, $\mathbf{g}_\alpha$ can be written as

$$\mathbf{g}_1 = \rho_{n1}\mathbf{v}_n + \rho_{s1}\mathbf{v}_{s1} + \tilde{\rho}(\mathbf{v}_{s1} - \mathbf{v}_{s2}) + \mathbf{J}^D \ , \tag{31}$$

$$\mathbf{g}_2 = \rho_{n2}\mathbf{v}_n + \rho_{s2}\mathbf{v}_{s2} + \tilde{\rho}(\mathbf{v}_{s2} - \mathbf{v}_{s1}) - \mathbf{J}^D, \tag{32}$$

where the normal density of the species α , $\rho_{n\alpha}$ is defined as

$$\rho_{n\alpha} \equiv \rho_\alpha - \rho_{s\alpha}, \qquad \rho_n = \rho_{n1} + \rho_{n2}. \tag{33}$$

From Eqs.(31) and (32), it is clear that the $\tilde{\rho}$ term describes the backflow created by one *superfluid* component on the other in the mixture.

Continuity equations and symmetry relations : The time evolution of the hydrodynamic variables $\{x_i\}$ are governed by continuity equations and conservation laws, which are

$$\partial_t\rho_\alpha = -\nabla\cdot\mathbf{g}_\alpha, \quad \alpha = 1, 2, \tag{34}$$

$$\partial_t\mathbf{g}_i = -\rho_1\nabla_i\phi_1 - \rho_2\nabla_i\phi_2 - \nabla_j\Pi_{ij}, \tag{35}$$

$$\partial_t\epsilon = -\nabla\cdot\mathbf{Q} - \mathbf{g}_1\cdot\nabla\phi_1 - \mathbf{g}_2\cdot\nabla\phi_2, \tag{36}$$

$$\partial_t\mathbf{v}_{s\alpha} = -\nabla\chi_\alpha, \quad \alpha = 1, 2, \tag{37}$$

where $\chi_\alpha = -\frac{\hbar}{m}\partial_t\theta_\alpha$, and ϕ_1 and ϕ_2 are the external potential experienced by Bosons **1** and **2** respectively, Eqs.(34) to (36) are continuity equations for density, momentum and energy respectively. Π_{ij} is the stress tensor and $\mathbf{Q}$ is the energy current. An argument of Martin *et al.,* shows that Π_{ij} can always be taken to be a symmetric tensor [10]. The quantities

$\{\mathbf{Q}, \Pi_{ij}, \mathbf{g}_\alpha, \chi_\alpha\} \equiv \{\mathcal{J}_i\}$, referred to as "currents", control the flow of the hydrodynamic variables $\{x_i\}$.

To complete the hydrodynamic description, we need to express the currents $\{\mathcal{J}_i\}$ in terms of the hydrodynamic variables $\{x_i\}$ so as to form a closed set of equations. As explained in the next section, this can be done by studying the entropy production of the system. The derivation can be simplified considerably by using the Galilean transformation properties of $\{\mathcal{J}_i\}$. By noting that the partial derivatives ∂_t and ∇ transform as $\partial_t \to \partial_{t'} = \partial_t - \mathbf{u}\cdot\nabla$, and $\nabla \to \nabla' = \nabla$ under a Galilean transformation, we have

$$Q'_i = Q_i + \Pi_{ij}u_j + u_i\epsilon' + \frac{1}{2}(g_{1i} + g_{2i})u^2 \tag{38}$$

$$\Pi'_{ij} = \Pi_{ij} + u_i(g_1 + g_2)_j + (g_1 + g_2)_i u_j + \rho u_i u_j \tag{39}$$

$$\chi'_\alpha = \chi_\alpha + \mathbf{u}\cdot\mathbf{v}_{s\alpha} + \frac{u^2}{2}. \tag{40}$$

For example, Eq.(40) is obtained by noting $\chi'_\alpha = -\frac{\hbar}{m_\alpha}(\partial_t - \mathbf{u}\cdot\nabla)(\theta_\alpha + m_\alpha\mathbf{u}\cdot\mathbf{r}/\hbar + m_\alpha u^2 t/2\hbar)$ $= \chi_\alpha - u^2/2 + \mathbf{u}\cdot\mathbf{v}_{s\alpha} + u^2$.

Using these properties, we can write Eqs.(38) to (40) as

$$Q_i = Q^0_i + \Pi^0_{ij}v_{nj} + v_{ni}\epsilon + \frac{1}{2}(g^0_1 + g^0_2)_i v^2_n, \tag{41}$$

$$\Pi_{ij} = \Pi^0_{ij} + v_{ni}(g^0_1 + g^0_2)_j + (g^0_1 + g^0_2)_i v_{nj} + \rho v_{ni}v_{nj}, \tag{42}$$

$$\chi_\alpha = \chi^0_\alpha + \mathbf{v}_n\cdot\mathbf{v}^0_{s\alpha} + \frac{1}{2}\mathbf{v}^2_n. \tag{43}$$

For later use, we note that

$$(\nabla\cdot\mathbf{g})^0 = \nabla\cdot\mathbf{g}^0 + \rho\nabla\cdot\mathbf{v}_n, \tag{44}$$

$$(\nabla\cdot\mathbf{Q})^0 = \nabla\cdot\mathbf{Q}^0 + \Pi^0_{ij}\partial_j v_{ni} + \epsilon^0(\nabla\cdot\mathbf{v}_n), \tag{45}$$

$$(\nabla\chi_\alpha)^0 = \nabla\chi^0_\alpha + (v^0_{s\alpha})_i\nabla v_{ni}. \tag{46}$$

II. Derivation of the hydrodynamic equations for binary mixtures of scalar superfluids :

According to the second law of thermodynamics, the entropy production of the system must be positive, $\frac{\mathrm{d}}{\mathrm{dt}}\int \mathrm{d}^3xs \geq 0$. Note that

$$\partial_t s = -\frac{1}{T}\nabla\cdot\mathbf{Q} + \frac{v_{ni}}{T}\partial_j\Pi_{ij} + \sum_{\alpha=1,2}\left(-\frac{1}{T}\mathbf{g}_\alpha\cdot\nabla\phi_\alpha + \frac{\mu_\alpha}{T}\nabla\cdot\mathbf{g}_\alpha + \frac{1}{T}\mathbf{b}_\alpha\cdot\nabla\chi_\alpha + \frac{1}{T}\rho_\alpha\mathbf{v_n}\cdot\nabla\phi_\alpha\right). \tag{47}$$

Evaluating Eq.(47) in the $\mathbf{v}_n = 0$ frame, we have

$$T(\partial_t s)^0 = -\left(\nabla \cdot \mathbf{Q}^0 + \Pi^0_{ij}\partial_i v_{nj} + \epsilon^0 \nabla \cdot \mathbf{v}_n\right) + \\ + \sum_{\alpha=1,2} [-\mathbf{g}^0_\alpha \cdot \nabla\phi_\alpha + \mu^0_\alpha \rho_\alpha \nabla \cdot \mathbf{v}_n + \mu^0_\alpha \nabla \cdot \mathbf{g}^0_\alpha + \mathbf{b}_\alpha \cdot \nabla \chi^0_\alpha \\ +(v^0_{s\alpha})_i (\mathbf{b}_\alpha \cdot \nabla) v_{ni}]. \tag{48}$$

Using Eq.(24), and Eqs.(44) to (46), Eq.(48) can be written as

$$T(\partial_t s)^0 = -\nabla \cdot [\mathbf{Q}^0 + \sum_{\alpha=1,2} (\mathbf{g}^0_\alpha \phi_\alpha - \mathbf{b}_\alpha \chi^0_\alpha)] + \\ -\left[\Pi^0_{ij} + (Ts - P)\delta_{ij} - \sum_\alpha (v^0_{s\alpha})_j b_{\alpha i}\right] \partial_i v_{nj} \\ + \sum_{\alpha=1,2} [(\phi_\alpha + \mu^0_\alpha)\nabla \cdot \mathbf{g}^0_\alpha - \chi^0_\alpha \nabla \cdot \mathbf{b}_\alpha]. \tag{49}$$

By noting that $\partial_t s + \nabla \cdot (s\mathbf{v}_n)= (\partial_t s)^0 + s\nabla \cdot \mathbf{v}_n$ is Galilean invariant and using Eq.(30), Eq.(49) can be written as

$$T[\partial_t s + \nabla \cdot (s\mathbf{v}_n)] = -\nabla \cdot \mathbf{Q}^D - \Pi^D_{ij}\partial_i v_{nj} - \sum_{\alpha=1,2} \chi^D_\alpha \nabla \cdot \mathbf{b}_\alpha + \nu \nabla \cdot \mathbf{J}^D, \tag{50}$$

where $\mathbf{J}^D$ is defined in Eq.(30). We introduce Π^D, $\mathbf{Q}^D$, χ^D, $\mathbf{J}^D$, and ν^D such that

$$\Pi^0_{ij} = \delta_{ij} P + \sum_{\alpha=1,2} b_{\alpha i} (v^0_{s\alpha})_j + \Pi^D_{ij}, \tag{51}$$

$$\mathbf{Q}^0 = \sum_{\alpha=1,2} \left(\mathbf{b}_\alpha \chi^0_\alpha - \phi_\alpha \mathbf{g}^0_\alpha\right) + \mathbf{Q}^D, \tag{52}$$

$$\chi^0_\alpha = \phi_\alpha + \mu^0_\alpha + \chi^D_\alpha, \tag{53}$$

$$\nu = (\mu^0_1 + \phi_1) - (\mu^0_2 + \phi_2). \tag{54}$$

The determination of the "currents" $\{\mathcal{J}_i\} = \{\mathbf{Q}, \Pi_{ij}, \mathbf{g}_\alpha, \chi_\alpha\}$ now reduces to the determination of their "dissipative" components $\{\mathcal{J}^D_i\} \equiv \{\mathbf{Q}^D, \Pi^D_{ij}, \mathbf{J}^D, \chi^D_\alpha\}$. Introducting R_1 and R_2 defined below, Eq.(47) becomes

$$\partial_t s + \nabla \cdot \left(s\mathbf{v}_n + \frac{\mathbf{Q}^D}{T} - \frac{\nu \mathbf{J}^D}{T}\right) = R \equiv R_1 + R_2, \tag{55}$$

$$R_1 = -\frac{1}{T}\left[\chi^D_1 \nabla \cdot \mathbf{b}_1 + \chi^D_2 \nabla \cdot \mathbf{b}_2 + \Pi^D_{ij}\partial_i v_{nj}\right] \ , \tag{56}$$

$$R_2 = \mathbf{Q}^D \cdot \nabla\left(\frac{1}{T}\right) - \mathbf{J}^D \cdot \nabla\left(\frac{\nu}{T}\right) = -(\mathbf{Q}^D - \nu \mathbf{J}^D) \cdot \left(\frac{\nabla T}{T^2}\right) - \mathbf{J}^D \cdot \left(\frac{\nabla \nu}{T}\right). \tag{57}$$

The right hand side of Eq.(55) expresses the entropy generated at $\mathbf{r}$ by the "external forces" $\{f_i\} \equiv \{-\nabla T, \partial_i v_{nj}, -\nabla \nu, \nabla \cdot \mathbf{b}_\alpha\}$. The dissipative "currents" $\{\mathcal{J}_i^D\}$ are responses to the disturbances $\{f_i\}$ as the system tries to relax to equilibrium. In the linear response limit, $\{\mathcal{J}_i^D\}$ is linear in $\{f_i\}$, and R becomes a quadratic form in the forces, which must be positive in order to satisfy the second law of thermodynamics. Once the relations between the dissipative currents $\{\mathcal{J}_i^D\}$ and the driving forces $\{f_i\}$ are determined, the hydrodynamic equations, given by Eqs.(34) to (37) will form a closed set. Using Eqs.(31) and (32) for $\mathbf{g}_\alpha$, Eqs.(42) and (51) for Π_{ij}, Eqs.(43) and (53) for χ_α, and Eqs.(41) and (52) for $\mathbf{Q}$, we can write Eqs.(34) to (37) as

$$\mathbf{I}. \qquad \partial_t \rho_\alpha = -\nabla \cdot \mathbf{g}_\alpha, \quad \alpha = 1,2, \tag{58}$$

$$\mathbf{g}_1 = \rho_{n1}\mathbf{v}_n + \rho_{s1}\mathbf{v}_{s1} + \tilde{\rho}\left(\mathbf{v}_{s1} - \mathbf{v}_{s2}\right) + \mathbf{J}^D, \tag{59}$$

$$\mathbf{g}_2 = \rho_{n2}\mathbf{v}_n + \rho_{s2}\mathbf{v}_{s2} + \tilde{\rho}\left(\mathbf{v}_{s2} - \mathbf{v}_{s1}\right) - \mathbf{J}^D, \tag{60}$$

$$\mathbf{II}. \qquad \partial_t(\mathbf{g}_1 + \mathbf{g}_2)_i = -\rho_1 \nabla_i \phi_1 - \rho_2 \nabla_i \phi_2 - \nabla_j (\Pi^r_{ij} + \Pi^D_{ij}), \tag{61}$$

$$\begin{aligned}\Pi^r_{ij} = \delta_{ij} P + (\rho_{n1} + \rho_{n2})\, v_{ni} v_{nj} + \textstyle\sum_{\alpha=1,2} \rho_{s\alpha} \left(v_{s\alpha}\right)_i \left(v_{s\alpha}\right)_j \\ + \tilde{\rho}\left(v_{s1} - v_{s2}\right)_i \left(v_{s1} - v_{s2}\right)_j ,\end{aligned} \tag{62}$$

$$\mathbf{III}. \quad \partial_t \mathbf{v}_{s\alpha} = -\nabla\left(\phi_\alpha + \mu_\alpha + \mathbf{v}_n \cdot \mathbf{v}_{s\alpha} + \chi^D\right), \quad \alpha = 1,2, \tag{63}$$

$$\mathbf{IV}. \quad \partial_t s + \nabla \cdot (s\mathbf{v}_n) + \nabla \cdot \left(\frac{\mathbf{Q}^D}{T} - \frac{\nu \mathbf{J}^D}{T}\right) = R, \tag{64}$$

where we have replaced the energy equation Eq.(36) by the entropy equation Eq.(64). For hydrodynamic processes with weak dissipation, the dissipative currents $\mathbf{J}^D, \Pi^D, \chi^D, \mathbf{Q}^D$ can be ignored and the resulting equations **I** to **IV** above will be referred to as "dissipation-free" hydrodynamic equations. For dissipative processes, however, it is necessary to determine the explicit form of the dissipative currents.

Determination of dissipative currents : Note that the forces in R_1 and R_2 (eq.(56) and Eq.(57)) have different parity and time reversal symmetries. As a result, the currents $(\chi_\alpha, \Pi^D_{ij})$ and $(\mathbf{Q}^D, \mathbf{J}^D)$ are only linear combinations of $(\nabla \cdot \mathbf{b}_\alpha, \partial_i v_{nj})$ and $(\nabla T, \nabla \nu)$ respectively. Since Π^D_{ij} is symmetric, it can be decomposed as

$$\Pi^D_{ij} = A_{ij} + \tau \delta_{ij}, \quad \mathrm{Tr} A = 0, \tag{65}$$

which allows R_1 to be expressed as

$$R_1 = -\frac{1}{T}\left[\frac{1}{2} A_{ij}\left(\nabla_i v_{nj} + \nabla_j v_{ni} - \frac{2}{3}\delta_{ij} \nabla \cdot \mathbf{v}_n\right) + \tau \nabla \cdot \mathbf{v}_n + \sum_{\alpha=1,2} \chi_\alpha (\nabla \cdot \mathbf{b}_\alpha)\right]. \tag{66}$$

Linear response implies

$$A_{ij} = -\eta\left(\nabla_i v_{nj} + \nabla_j v_{ni} - \frac{2}{3}\delta_{ij} \nabla \cdot \mathbf{v}_n\right), \tag{67}$$

$$\begin{pmatrix} \tau \\ \chi_1^D \\ \chi_2^D \end{pmatrix} = - \begin{pmatrix} \zeta_{00} & \zeta_{01} & \zeta_{02} \\ \zeta_{01} & \zeta_{11} & \zeta_{12} \\ \zeta_{02} & \zeta_{21} & \zeta_{22} \end{pmatrix} \begin{pmatrix} \nabla \cdot \mathbf{v}_n \\ \nabla \cdot \mathbf{b}_1 \\ \nabla \cdot \mathbf{b}_2 \end{pmatrix}, \tag{68}$$

where η will be referred to as the "first" viscosity, and ζ_{ij} as the "second" viscosity matrix. The fact that ζ_{ij} is symmetric is a consequence of the general Onsager relations for kinetic coefficients. More explicitly, we have

$$\Pi_{ij}^D = -\eta \left(\nabla_i v_{nj} + \nabla_j v_{ni} - \frac{2}{3} \delta_{ij} \nabla \cdot \mathbf{v}_n \right) - \zeta_{00} \nabla \cdot \mathbf{v}_n - \zeta_{01} \nabla \cdot \mathbf{b}_1 - \zeta_{02} \nabla \cdot \mathbf{b}_2, \tag{69}$$

and

$$\chi_\alpha^D = -\zeta_{0\alpha} \nabla \cdot \mathbf{v}_n - \sum_{\sigma=1,2} \zeta_{\alpha\sigma} \nabla \cdot \mathbf{b}_\sigma. \tag{70}$$

Recall from Eq.(27) that $\mathbf{b}_1 = \rho_{s1} \mathbf{v}_{s1}^0 + \tilde{\rho}(\mathbf{v}_{s1}^0 - \mathbf{v}_{s2}^0)$ and $\mathbf{b}_2 = \rho_{s2} \mathbf{v}_{s2}^0 + \tilde{\rho}(\mathbf{v}_{s2}^0 - \mathbf{v}_{s1}^0)$. Clearly, in order to have $R_1 \geq 0$, we need to have $\eta > 0$, and $\det|\zeta| \geq 0$.

To find the dissipative currents $\mathbf{Q}^D$ and $\mathbf{J}^D$, we note that the expression for R_2 is identical to the entropy production of a classical binary mixture [11]. The expressions of these currents are therefore identical to those of classical fluids and are of the form

$$\mathbf{J}^D = -\alpha T \left(\frac{\nabla \nu}{T} \right) - \beta T^2 \left(\frac{\nabla T}{T^2} \right) \tag{71}$$

$$\mathbf{Q}^D - \nu \mathbf{J}^D = -\beta T^2 \left(\frac{\nabla \nu}{T} \right) - \gamma T^2 \left(\frac{\nabla T}{T^2} \right), \tag{72}$$

where we have again used the Onsager relation to identify the off-diagonal kinetic coefficients in Eqs.(71) and (72). Eq.(57) then becomes $R_2 = \gamma (\nabla T)^2/T^2 + \alpha (\nabla \nu)^2/T + 2\beta \left(\nabla T \cdot \nabla \nu \right)/T$, or,

$$R_2 = \kappa \frac{(\nabla T)^2}{T^2} + \frac{1}{\alpha T} \left(\mathbf{J}^D \right)^2, \tag{73}$$

where $\kappa \equiv \gamma - \beta^2 T/\alpha$. The positivity of R_2 implies $\kappa \geq 0$, and $\alpha \geq 0$. Eqs.(69), (70), (71), and (72) completely determine the dissipative currents and complete the full dissipative hydrodynamic equations, given by Eqs.(58) to (64) together with Eqs.(69) to and Eq.(72).

While we shall not be solving the hydrodynamic equations here, we remind the readers that the sound modes of the system are contained in the so-called linearized hydrodynamic equations, which are obtained by linearizing Eqs.(58) to (64) about the equilibrium configuration. In the single component superfluid case, Landau [6] has pointed out that it is useful to express all the hydrodynamic variables in terms of externally controllable quantities such as temperature T and pressure P. (Also see reference [8]). In the mixture case, there is another controllable quantity, which is the concentration $c = \rho_1/\rho$ defined earlier in Section **I**. Expressing $\mathbf{J}^D$ and $\mathbf{Q}^D$ in terms of T, P and c as in Landau and Lifshitz [11], we have

$$\mathbf{J}^D = -\rho D \left(\nabla c + \frac{k_T}{T} \nabla T + \frac{k_P}{P} \nabla p \right), \tag{74}$$

$$\mathbf{Q}^D = \left[k_T \left(\frac{\partial \nu}{\partial c} \right)_{P,T} - T \left(\frac{\partial \nu}{\partial T} \right)_{P,c} + \nu \right] \mathbf{J}^D - \kappa \nabla T, \tag{75}$$

where κ, D, k_T, k_P are the thermal conductivity, the diffusion coefficient, the coefficient of thermal diffusion, and the coefficient of baro-diffusion respectively; they can be written as

$$\kappa = \gamma - \beta^2 T/\alpha, \qquad D = \frac{\alpha}{\rho} \left(\frac{\partial \nu}{\partial c} \right)_{T,P} \tag{76}$$

$$\rho D k_T/T = \alpha \left(\frac{\partial \nu}{\partial T} \right)_{P,c} + \beta, \qquad \rho D k_P/P = \alpha \left(\frac{\partial \nu}{\partial P} \right)_{T,c}. \tag{77}$$

Using the relations above, the full expression for entropy production is

$$\begin{aligned} R = & \frac{\eta}{2T} \left(\nabla_i v_{nj} + \nabla_j v_{ni} - \frac{2}{3} \delta_{ij} \nabla \cdot \mathbf{v}_n \right)^2 \\ & + \frac{\zeta_{00}}{T} \left(\nabla \cdot \mathbf{v}_n \right)^2 + 2 \sum_{\alpha=1,2} \frac{\zeta_{0\alpha}}{T} \left(\nabla \cdot \mathbf{v}_n \right) \left(\nabla \cdot \mathbf{b}_\alpha \right) \\ & + \sum_{\alpha=1,2} \frac{\zeta_{\alpha\alpha}}{T} \left(\nabla \cdot \mathbf{b}_\alpha \right)^2 + +2 \sum_{\alpha,\sigma=1,2} \frac{\zeta_{\alpha\sigma}}{T} \left(\nabla \cdot \mathbf{b}_\alpha \right) \left(\nabla \cdot \mathbf{b}_\sigma \right) \qquad (78) \\ & + \kappa \frac{(\nabla T)^2}{T^2} + \frac{D\rho}{T} \frac{\partial \nu}{\partial c} \left(\nabla c + \frac{\kappa_T}{T} \nabla T + \frac{\kappa_P}{P} \nabla p \right)^2. \qquad (79) \end{aligned}$$

III. Hydrodynamic equations of other mixture systems :
(a) Binary mixture of a normal fluid and a superfluid : Let component **1** be the superfluid and component **2** be the normal fluid. Since **2** does not have broken gauge symmetry, $\mathbf{v}_{s2}$ is absent in the hydrodynamic equations. Rotational symmetry (Eq.(20)) yeilds $\mathbf{b}_1 \times \mathbf{v}^o_{s1} = 0$, which implies $\mathbf{b}_1 = \rho_{s1} \mathbf{v}_{s1}$, or $\rho_2 = \tilde{\rho} = 0$. Following exactly the same steps as before, it is easy to see that the the hydrodynamic equations of this system are still given by Eqs.(58) to (64) with the quantities $(\rho_{s2}, \tilde{\rho}, \mathbf{v}_{s2})$ all set to zero. Moreover, the dissipative part of the stress tensor Π^D_{ij} as well as χ^D_1 are given by Eqs.(69) and (70) with ζ_{02} and $\mathbf{b}_2$ set to zero, while the expressions of the diffusion current $\mathbf{J}^D$ and the heat current $\mathbf{Q}^D$ are still given by Eqs.(74) and (75).

(b) Binary mixture of normal fluids :

In this case, both $\mathbf{v}_{s1}$ and $\mathbf{v}_{s2}$ are absent from the equations. The hydrodynamic equations are given by Eqs.(58) to Eq.(64) with the quantities $(\rho_{s1}, \rho_{s2}, \tilde{\rho}, \mathbf{v}_{s1}, \mathbf{v}_{s2})$ set to zero. The dissipative component Π^D is given by Eq.(69) with all $\zeta_{\alpha\beta} = 0$ except for ζ_{00}, and $\chi^D = 0$. The diffusion current and heat current are still given by Eqs.(74) and (75).

Acknowledgement : This work is support in part by NSF Grant DMR-9705295.

REFERENCES

[1] M.H. Anderson, J.R. Ensher, M.R. Mathews, C.E. Weiman, and E.A. Cornell, Science **269**, 198 (1995).

[2] K.B. Davis, M.O. Mewes, M.R. Andrews, N.J. van Druten, D.S. Durfee, D.M. Kurn, and W. Ketterle, Phys. Rev. Lett. **75**, 3969 (1995).

[3] C.C. Bradley, C.A. Sackett, J.J. Tollett, and R.G. Hulet, Phys. Rev. Lett. **75**, 1687 (1995).

[4] Tin-Lun Ho and V.B. Shenoy, Phys. Rev. Lett. **77**, 3276 (1996).

[5] C.J. Myatt, E. Cornell, and C. Wieman, Phys. Rev. Lett. **78**, 586 (1997).

[6] L.D. Landau, J. Phys (USSR) **5**, 71 (1941); L.D. Landau and E.M. Lifshitz, *Fluid Mechanics* (Pergamon Press, New York, 1987), 2nd ed., Chapter XVI.

[7] The method used here follows closely to the one constructed by one of us for the hydrodynamics of superfluid ^{3}He-A, Tin-Lun Ho in *Sanibel Symposium on Quantum Fluids and Solids*, edited by S.B.Trickey, E.D.Adams, and J.W.Duffy, p.97, (1977).

[8] V.B. Shenoy and T.L.Ho, to be published.

[9] The spinor effects are discussed in Tin-Lun Ho and V.B. Shenoy, Phys. Rev. Lett. **77**, 2595 (1996).

[10] See the Appendix of P.C. Martin, O. Parodi, and P.S. Pershan, Phys. Rev. **A**6, 2401 (1972).

[11] See L.D. Landau and E.M. Lifshitz, *Fluid Mechanics* (Pergamon Press, New York, 1987), 2nd ed., Chapter VI.